



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

163. Proposed by R. D. CARMICHAEL, Anniston, Ala.

In a regular n -gon a triangle is formed by taking three vertices at random. What is the mean value of the triangle?

No solution has been received.

164. Proposed by J. O. MAHONEY, B. E., M. Sc., Central High School, Dallas, Tex.

If m is prime, and the numbers $0, 1, 2, \dots, m^2 - 1$ are placed at random in the form of a square, the probability that the square is hyper-magic is $(m-1)m/(m^2-2)!$

No solution has been received.

167. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

A line l is divided into n segments by $n-1$ points taken at random on it; find the mean value of the product of p of the segments, the p segments being taken at random and p being less than n .

Solution by the PROPOSER.

Let $a, b, c, \dots$ be the p segments in some particular case. The chance that another point taken at random shall fall on the first of these segments is a/l ; on the second, b/l ; and so on. Hence the chance in this particular case that p new points taken at random will fall each on one of the p segments $a, b, c, \dots$ in an assigned order is $abc\dots/l^p$. The chance that they shall so fall in any order is evidently $p!abc\dots/l^p$. Hence the probability of this occurring, however the line is divided and however the p points are chosen, is $p!m(abc\dots)/l^p$.

If now we can find another expression involving no unknown quantity and giving this same probability, the two will enable us to determine the value of $m(abc\dots)$.

The number of ways in which p points can be taken one on each of p chosen segments is the same as the number of ways in which $p-1$ points can be placed one between the two of each consecutive pair of p other points, and is easily found by the theory of permutations to be $p!(p-1)!$. But p segments may be chosen from n segments in $\frac{n!}{p!(n-p)!}$ ways. Hence the whole number of ways in which p points can be chosen on n segments one on each of p segments is

$$\frac{p!(p-1)!n!}{p!(n-p)!} = \frac{n!(p-1)!}{(n-p)!}.$$

Now the whole number of ways in which $n-1+p$ points can be arranged is $(n+p-1)!$. Hence the chance that no two of the p points last chosen shall be on the same one of the original n segments is

$$\frac{n!(p-1)!}{(n-p)!} \div (n+p-1)! = \frac{n!(p-1)!}{(n-p)!(n+p-1)!}.$$